



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

an hour in the dense chapparral below, but found nothing. Probably, the shot being small, it was but slightly hurt. Neither of us had ever seen this species before in the wild state, although Mr. Hawley is a native of Southern California and has been often in these mountains.

In this connection I beg to correct a mistake in my note on *Fregata aquila*, published in 'The Auk' for October, 1893 (p. 362). In my reference to the Humboldt Bay specimen, I should have quoted Mr. T. S. Palmer instead of Mr. Anthony as the authority for the record. — R. H. LAWRENCE, *Monrovia, Cal.*

Capture of the Golden Eagle at Covington, Virginia. — It gives me pleasure to record the capture of a fine male specimen of the Golden Eagle (*Aquila chrysaëtos*) at Covington, Alleghany County, Virginia, on Oct. 28, 1893. It was shot by Derry B. Smith, Esq., who kindly sent me the specimen for mounting. On skinning I found it very fat. It measured as follows: Length, 33 inches; extent, 78 inches; wing, 24 inches; tail, 14 inches. This is the first specimen, to my knowledge, that has ever been taken in this immediate vicinity. — THADDEUS SURBER, *White Sulphur Springs, W. Va.*

Another Record of the Breeding of the Saw-whet Owl (*Nyctale acadica*) in Eastern Massachusetts. — As there are still but few records of the breeding of the Saw-whet Owl in eastern Massachusetts, I take pleasure in adding one more.

On July 3, 1893, Mr. Gerrit S. Miller, Jr., and I were setting a line of traps in a heavy white pine swamp that lies along Red Brook in the town of Wareham, Mass. We noticed a large old pine stump which was broken off at about 25 feet above the ground and full of Woodpeckers' holes, and pounded on it. We had pounded but once or twice when a Saw-whet Owl popped her head out of the uppermost hole and kept it there motionless, although I fired at her three times with my pistol. The third shot killed her and she fell back into the hole.

On taking the bird out, I found there was a nest containing seven eggs. The nest was quite bulky and composed of gray moss (*Usnea*) interwoven with small pieces of fibrous bark, a few pine needles, small twigs, and feathers of the bird herself. The hole in which the nest was found was 18 feet from the ground and about 8 inches deep.

In the nest besides the eggs was a half eaten red-backed mouse (*Evo-
tomys gapperi*).

Three of the eggs were in various stages of incubation, one being on the point of hatching, — in fact the young bird had already cracked the shell. Three were addled, and one was perfectly fresh.

On dissecting the old bird we found that she had laid her full set of eggs. Her stomach contained the other half of the *Evo-
tomys*, which she was apparently eating when we disturbed her.

I believe the only other records for Massachusetts are:—

'Probable breeding of the Acadian Owl (*Nyctale acadica*) in Massachusetts.' R. Deane, Bull. Nutt. Orn. Club, Vol. II, July, 1877, p. 84. Three specimens in first plumage are recorded,—one, taken June 28, 1876, at Newton, Mass., one at Hingham, Mass., July 5, 1876, and one July 8, 1876.

'Breeding of the Acadian Owl in Eastern Massachusetts.' N. A. Francis Bull. Nutt. Orn. Club, Vol. VI, July, 1881, p. 185. Nest with five young found June 4, 1880, at Braintree, Mass.

'Breeding of the Acadian Owl (*Nyctale acadica*) in Massachusetts.' Bull. Nutt. Orn. Club, Vol. VI, July, 1881, pp. 143-145. Account by William Brewster of nest with four eggs taken at 'Tyngsboro', Mass., April 5, 1881, by W. B. Perham. *Ibid.*, Jan., 1882, pp. 23-25. Additional notes on nesting at Tyngsboro', by W. B. Perham, who found seven nests in all.

'Ornithologist and Oologist,' Vol. XIV, Oct., 1889, pp. 155-156. Record of nest with four eggs, well advanced in incubation, taken at Dunstable, Mass., May 1, 1889, by C. W. Swallow.

In connection with this see also account of four nests found at Holland Patent, N. Y., by Egbert Baggs, in 'Ornithologist and Oologist,' Vol. XII, No. 4, April, 1887, p. 57.—OUTRAM BANGS, *Wareham, Mass.*

Capture of Another Flammulated Owl in California.—On May 26, 1893, I became the possessor of an Owl which after a careful examination Mr. F. Stephens decides is *Megascops flammeola*. As this is only the *fourth* specimen known to have been taken in this State, I thought it might be of some interest to the readers of 'The Auk' to know of it. This specimen was taken in the San Bernardino range of mountains at an elevation of 5000 feet. The specimen was a male and measured as follows: Length, 7.50 inches; alar extent, 17.50.—E. D. PALMER, *San Bernardino, Cal.*

Empidonax flaviventris on Long Island, N. Y.—While collecting at Flatbush, in the suburbs of Brooklyn, on June 4, 1892, I secured a male Yellow-bellied Flycatcher, which I find is the first record for Long Island. No others were observed, although I hunted carefully through the patch of underbrush and dead saplings where the specimen was secured.—CURTIS CLAY YOUNG, *Brooklyn, N. Y.*

Corrections.—*Xanthocephalus xanthocephalus* and *Spiza americana* in Maine.—In a note in 'The Auk,' Vol. X, July, 1893, p. 302, I mentioned these birds in terms that require further notice. The specimen of *X. xanthocephalus* brought under consideration was first noted by Mr. Ridgway in 1887 (*cf.* Auk, Vol. IV, July 1887, p. 256). But in the dates given in the two notices there is a discrepancy of nearly a year. As I saw the bird before and at the time it was shot, and kept record of the fact in my diary, I feel authorized to furnish the correct date, which is, as I have previously stated, Aug. 17, 1882. That both notes refer to the